

# Population Practice Profile – page 1

DQCMSv2.1 BETA

## Population Practice Profile CLINIC TEST

Criteria Selected:  
None selected

Total number of active patients for the criteria	No. of Patients	
10		
<b>Office Visits</b>	<b>%</b>	<b>(#)</b>
Patients with office visits from 03/12/08 to 03/12/09	50.0	(5 / 10)
<b>A1c</b>		
Patients who have had an A1C from 03/12/08 to 03/12/09	30.0	(3 / 10)
Of patients with A1C from 03/12/08 to 03/12/09, the percent whose most recent is less than 7 %	0.0	(0 / 3)
Patients who have had 2 or more A1c results separated by more than 3 months in the past 12 months	0.0	(0 / 10)
<b>Lipid Profile</b>		
Patients >= 18 who have had an LDL from 03/12/08 to 03/12/09	22.2	(2 / 9)
Of patients with LDL from 03/12/08 to 03/12/09, the percent whose most recent LDL is < 100 mg/dl	50.0	(1 / 2)
Of patients with most recent LDL >= 100 mg/dl, the percent having lipid lowering medication prescribed	100.0	(4 / 4)
Of patients with HDL from 03/12/08 to 03/12/09, the percent whose most recent HDL is > 45 mg/dl	100.0	(2 / 2)
Of patients with triglyceride from 03/12/08 to 03/12/09, the percent whose most recent triglyceride is < 150 mg/dl	50.0	(1 / 2)
<b>Kidney</b>		
Patients who have had a creatinine test from 03/12/08 to 03/12/09	20.0	(2 / 10)
Patients who have had an e-GFR test from 03/12/08 to 03/12/09	20.0	(2 / 10)
Patients who have had an albuminuria test from 03/12/08 to 03/12/09	20.0	(2 / 10)
Of patients with "+" albuminuria ever, the percent having ACE-I / ARB prescribed	66.7	(2 / 3)
<b>Blood Pressure</b>		
Patients with diagnosed HTN	20.0	(2 / 10)
Patients with Dx=HTN, the percent having ACE-I / ARB prescribed	100.0	(2 / 2)
Patients with Dx=HTN with microalbumin test from 03/12/08 to 03/12/09	0.0	(0 / 1)
Patients with Dx=HTN who have ever had positive urine protein or microalbumin, the percent having ACE-I / ARB prescribed	100.0	(1 / 1)
Of patients with Dx=HTN, the percent whose average of the three most recent systolic < 130 and diastolic < 80	50.0	(1 / 2)
Of patients with Dx=HTN whose average of the three most recent systolic >= 130 or diastolic >= 80, percent who have ACE-I / ARB prescribed	100.0	(1 / 1)
Of patients without Dx=HTN, the percent whose average of the three most recent systolic < 130 and diastolic < 80	60.0	(3 / 5)

Date Generated: 03/12/09

# Population Practice Profile – page 2

DQCMSv2.1 BETA

## Population Practice Profile CLINIC TEST

Criteria Selected:  
None selected

Total number of active patients for the criteria	No. of Patients	
10		
<b>Foot Exam</b>	<b>%</b>	<b>#</b>
Patients who have had any foot exam from 03/12/08 to 03/12/09	30.0	(3 / 10)
Patients who have had a monofilament foot exam from 03/12/08 to 03/12/09	0.0	(0 / 10)
Of patients categorized as high risk ever, the percent who have had a monofilament foot exam from 03/12/08 to 03/12/09	0.0	(0 / 3)
Of patients categorized as high risk ever, those who have received referral to a foot care specialist	33.3	(1 / 3)
Of patients categorized as high risk ever, those who have received referral for therapeutic shoes	0.0	(0 / 3)
Patients who have had any foot exam from 03/12/08 to 03/12/09, the percent who have had an active foot problem	0.0	(0 / 3)
Of patients identified with an active foot problem within the last 3 months, the percent with follow-up office visit	0.0	(0 / 0)
<b>Dilated Eye / Dental Exam</b>		
Patients with a dilated eye exam from 03/12/08 to 03/12/09	20.0	(2 / 10)
Of patients who had any retinopathy ever, the percent with dilated eye exam from 03/12/08 to 03/12/09	66.7	(2 / 3)
Patients referred for a dilated eye exam from 03/12/08 to 03/12/09	0.0	(0 / 8)
Patients referred for a dental exam from 03/12/08 to 03/12/09	20.0	(2 / 10)
<b>Prevention Services</b>		
Patients who have ever had a pneumococcal vaccine	30.0	(3 / 10)
Patients who have had a influenza vaccine from 09/01/08 to 03/31/09	0.0	(0 / 10)
Patients $\geq$ 30 who have ASA / Anti-coagulation therapy prescribed	66.7	(6 / 9)
<b>Tobacco Use</b>		
Patients currently smoking/ using tobacco	60.0	(3 / 5)
Of those patients currently smoking/ using tobacco, the percent receiving cessation intervention from 03/12/08 to 03/12/09	100.0	(3 / 3)
Of patients currently smoking/using tobacco, percent participating in/using	0.0	(0 / 3)

Date Generated: 03/12/09